

Impaired microRNA processing causes corpus luteum insufficiency and infertility in mice

Motoyuki Otsuka, ... , Shengcai Lin, Jiahuai Han

J Clin Invest. 2008;118(6):2366-2366. <https://doi.org/10.1172/JCI33680E1>.

Erratum

Original citation: *J. Clin. Invest.* 118:1944–1954 (2008). doi:10.1172/JCI33680. Citation for this erratum: *J. Clin. Invest.* 118:2366 (2008). doi:10.1172/JCI33680E1. During the preparation of the manuscript, panels D and E in Figure 5 were inadvertently mislabeled. The correctly labeled panels and their corresponding legends appear below. The JCI regrets the error.

Find the latest version:

<https://jci.me/33680E1/pdf>





Erratum

Impaired microRNA processing causes corpus luteum insufficiency and infertility in mice

Motoyuki Otsuka, Min Zheng, Masaaki Hayashi, Jing-Dwan Lee, Osamu Yoshino, Shengcai Lin, and Jiahuai Han

Original citation: *J. Clin. Invest.* **118**:1944–1954 (2008). doi:10.1172/JCI33680.

Citation for this erratum: *J. Clin. Invest.* **118**:2366 (2008). doi:10.1172/JCI33680E1.

During the preparation of the manuscript, panels D and E in Figure 5 were inadvertently mislabeled. The correctly labeled panels and their corresponding legends appear below.

(D) miR17-5p and let7b had no effect on the proliferation of SVEC. Thirty-six hours after transfection of the indicated oligos into SVEC, 5×10^4 cells were seeded, and the number of cells was determined by cell counting at the indicated time points. Data are presented as mean \pm SD ($n = 4$ per group, $P = NS$). **(E)** miR17-5p and let7b had no effect on the motility of SVEC. In vitro wound healing was quantified as the average length of the elongation of wound edges at 4 specific points over 24 hours. Data are presented as mean \pm SD ($n = 3$ per group, $P = NS$).

The JCI regrets the error.

